

Unclassified

AD-A257 404

SECURITY CLASSIFICATION OF THIS PAGE

| | | | | | | |
|--|-------|--|---|---|--------------------------------|----------------------|
| REPORT DOCUMENTATION DTIC SELECT OCT 27 1992 S B | | | AD-A257 404 [Barcode] | | m Approved 18 No. 0704-0188 | |
| 1a. REPORT SECURITY CLASSIFICATION Unclassified | | | 1d. RESTRICTIVE MARKINGS | | | |
| 2a. SECURITY CLASSIFICATION AUTHORITY OCT 27 1992 | | | 3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for Public Release Distribution Unlimited | | | |
| 2b. DECLASSIFICATION / DOWNGRADING SCHEDULE B | | | 4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. DU/DC/TR-31 | | | |
| 6a. NAME OF PERFORMING ORGANIZATION Department of Chemistry Duke University | | 6b. OFFICE SYMBOL (If applicable) | | 7a. NAME OF MONITORING ORGANIZATION Office of Naval Research | | |
| 6c. ADDRESS (City, State, and ZIP Code) Durham, NC 27708-0346 | | | 7b. ADDRESS (City, State, and ZIP Code) 300 North Quincy Street Arlington, VA 22217-5000 | | | |
| 8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research | | 8b. OFFICE SYMBOL (If applicable) | | 9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-89-J1545 R&T Code 4135008 | | |
| 8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217-5000 | | | 10. SOURCE OF FUNDING NUMBERS | | | |
| | | | PROGRAM ELEMENT NO. NR | | PROJECT NO. 053 | TASK NO. 841 |
| | | | WORK UNIT ACCESSION NO. | | | |
| 11. TITLE (Include Security Classification) X-Ray Crystal Structure of an Organoindium Alkoxide Dimer, (Ph ₂ InOSiMe ₃) ₂ . | | | | | | |
| 12. PERSONAL AUTHOR(S) Mark F. Self, Andrew T. McPhail, and Richard L. Wells* | | | | | | |
| 13a. TYPE OF REPORT Technical | | 13b. TIME COVERED FROM _____ TO _____ | | 14. DATE OF REPORT (Year, Month, Day) 1992-10-20 | | 15. PAGE COUNT 12 |
| 16. SUPPLEMENTARY NOTATION Accepted for Publication in <i>J. Coord. Chem.</i> | | | | | | |
| 17. COSATI CODES | | | 18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) | | | |
| FIELD | GROUP | SUB-GROUP | indium-oxygen, alkoxide, ring compound, synthesis, crystal structure | | | |
| | | | | | | |
| 19. ABSTRACT (Continue on reverse if necessary and identify by block number) The solid-state structure of (Ph ₂ InOSiMe ₃) ₂ 1 has been established by single-crystal X-ray analysis. Monoclinic crystals of 1 belong to space group <i>P</i> 2 ₁ / <i>c</i> , with <i>a</i> = 9.136(1), <i>b</i> = 15.503(2), <i>c</i> = 11.836(1) Å, β = 102.46(1)°, <i>Z</i> = 2. Refinement of atomic parameters converged at <i>R</i> = 0.050 (<i>R</i> _w = 0.063) over 1718 observed reflections with <i>I</i> > 3.0σ(<i>I</i>). The dimeric molecule lies on a crystallographic centre of symmetry with In-O distance of 2.15(7) and 2.149(8) Å, O-In-O = 78.9(3)°, and In-O-In = 101.1(3)°. Compound 1 is only the second example of a simple dimeric organoindium alkoxide to be structurally characterized in this manner. | | | | | | |
| 20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS | | | 21. ABSTRACT SECURITY CLASSIFICATION Unclassified | | | |
| 22a. NAME OF RESPONSIBLE INDIVIDUAL Richard L. Wells | | | 22b. TELEPHONE (Include Area Code) (919) 660-1541 | | 22c. OFFICE SYMBOL | |

OFFICE OF NAVAL RESEARCH

Grant NO0014-89-J-1545

R&T Code 4135008

Technical Report No. DU/DC/TR-31

X-RAY CRYSTAL STRUCTURE OF AN ORGANOINDIUM ALKOXIDE DIMER, $(\text{Ph}_2\text{InOSiMe}_3)_2$

by

Mark F. Self, Andrew T. McPhail, and Richard L. Wells*

Prepared for Publication in *J. Coord. Chem.*

Duke University
Department of Chemistry,
P. M. Gross Chemical Laboratory
Box 90346
Durham, NC 27708-0346

October 20, 1992

118225
92-28189



15P6

92 10 26 220

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

(Submitted to *J. Coord. Chem.* 26 June 1992)
(Revised submitted 6 October 1992)

NOTE

**X-RAY CRYSTAL STRUCTURE OF AN ORGANOINDIUM
ALKOXIDE DIMER, $(\text{Ph}_2\text{InOSiMe}_3)_2$**

Mark F. Self, Andrew T. McPhail and Richard L. Wells *

*Department of Chemistry, Paul M. Gross Chemical Laboratory, Duke University,
Durham, North Carolina 27706, U S A.*

(Received, 1992)

The solid-state structure of $(\text{Ph}_2\text{InOSiMe}_3)_2$ **1** has been established by single-crystal X-ray analysis. Monoclinic crystals of **1** belong to space group $P2_1/c$, with $a = 9.136(1)$, $b = 15.503(2)$, $c = 11.836(1)$ Å, $\beta = 102.46(1)^\circ$, $Z = 2$. Refinement of atomic parameters converged at $R = 0.050$ ($R_w = 0.063$) over 1718 observed reflections with $I > 3.0\sigma(I)$. The dimeric molecule lies on a crystallographic centre of symmetry with In-O = 2.159(7) and 2.149(8) Å, O-In-O = $78.9(3)^\circ$, and In-O-In = $101.1(3)^\circ$. Compound **1** is only the second example of a simple dimeric organoindium alkoxide to be structurally characterized in this manner.

Keywords: indium, alkoxide, siloxide, X-ray structure

INTRODUCTION

Chemical investigations involving alkoxides of indium have recently undergone a resurgence^{1,2} due in part to their potential as precursors to III-VI materials some of which have found utility in applications such as conducting and transparent oxide films for displays and solar cell windows.³ Although a variety of organoindium alkoxides have been fabricated for use in relation to these

* Author for correspondence

DTIC QUALITY INSPECTED 2

| | |
|--------------------|-------------------------|
| By _____ | |
| Distribution/ | |
| Availability Codes | |
| Dist | Avail and/or Special |
| A-1 | |

material science studies, determinations of their detailed structures have thus far been lacking.

During the period 1956 to 1974, there appeared numerous literature reports of simple organoindium alkoxides (*e.g.*, R_2InOR' : $R = Me$; $R' = SiMe_3$,⁴ $SiPh_3$,⁵ CMe_3 ,⁶ CPh_3 ,⁷ Me ,⁸ CD_3 ,⁹ $t-Bu$ ¹⁰: $R = CD_3$; $R' = Me$, CD_3 ,⁹ $R = Et$; $R' = Me$, Et ⁹ $R = Bu$; $R' = Bu$, Ph ¹¹). The nature of these compounds was established by a combination of NMR spectroscopy, cryoscopic molecular weight determination, and infrared techniques. It is surprising that no X-ray crystallographic analyses of these compounds have appeared to date. In fact, the first and only report of the determination of the crystal structure of a simple dimeric organoindium alkoxide, $(t-Bu_2InOEt)_2$, was published only recently.¹² The solid-state structure of $(Ph_2InOSiMe_3)_2$ described herein represents only the second complete characterization of a member of this class.

EXPERIMENTAL

Materials and Measurements

All manipulations were performed using general Schlenk techniques in a Vacuum/Atmospheres HE-493 Dri-Lab containing an argon atmosphere. Toluene and benzene- d_6 were distilled from sodium/benzophenone ketyl under dry dinitrogen. $P(SiMe_3)_3$ ¹³ and Ph_2InCl ¹⁴ were prepared by literature procedures. The 1H NMR spectrum was recorded on a Varian XL-300 spectrometer using a flame sealed 5 mm tube and referenced to TMS by use of the residual protons of deuterated benzene at δ 7.15 ppm. The melting point (Thomas-Hoover Uni-melt, sealed capillary) is uncorrected. Elemental analysis was performed by E+R Microanalytical Laboratory, Inc., Corona, New York.

Preparation of (Ph₂InOSiMe₃)₂ 1

The title compound was prepared *via* the adventitious hydrolysis of a reaction mixture containing P(SiMe₃)₃ and Ph₂InCl in 100 mL of toluene. This hydrolysis occurred through the probable contamination of the solvent by a trace amount of water. The procedure which was employed follows. In the dry box, Ph₂InCl (0.250 g, 0.82 mmol) in 25 mL of toluene and P(SiMe₃)₃ (0.103 g, 0.41 mmol) in 75 mL of toluene were combined in a 150-mL one-necked round-bottomed flask equipped with a Teflon valve and a micro-stirbar. The resulting *milky* suspension was taken out of the box and stirred at room temperature for 24 h. Removal of the volatiles *in vacuo* yielded an oily off-white residue that was dissolved in approximately 15 mL of hot toluene and cooled to -15 °C for 1 day. Compound 1 was isolated in 71% yield (0.416g, based on Ph₂InCl) from an intractable yellow oil as clear colourless platelets. Mp = 230 °C (dec), Anal. Calcd (Found) for C₃₀H₃₈In₂O₂Si₂: C, 50.30 (49.93); H, 5.35 (5.46). ¹H NMR: δ 0.02 (s, 18 H, OSiMe₃), 7.02 (m, 20 H, Ph).

X-Ray Crystal Structure Analysis of Compound 1

All measurements were made on a sample mounted inside a flame-sealed 0.6 mm thin-walled glass capillary under an inert argon atmosphere. Oscillation and Weissenberg photographs provided preliminary unit cell parameters and space group information. Crystallographic data are summarized in Table I. The monoclinic space group *P*2₁/*c* was established unequivocally from the Laue symmetry and systematic absences (*0k0* when *k* ≠ 2*n*, *h0l* when *l* ≠ 2*n*). Intensity data (+*h*, +*k*, ±*l*, 3590 reflections, θ_{max} = 75°), recorded on an Enraf-Nonius CAD-4 diffractometer using Cu-Kα radiation (λ = 1.5418 Å, graphite monochromator), were corrected for the usual Lorentz and polarization effects. An empirical absorption correction, based on the φ-dependency of the intensities of two reflections with ψ ca. 90°, was applied, and equivalent reflections were averaged

[$R_{\text{merge}} = 0.038$ on (I)] to yield 3378 reflections out of which those 1718 with $I > 3.0\sigma(I)$ were retained for the analysis.

The crystal structure was solved by the heavy-atom approach. Approximate indium atom coordinates were derived from a Patterson map. The remaining non-hydrogen atoms were located in a weighted F_o Fourier synthesis phased by the indium atom. Several rounds of full-matrix least-squares adjustment of non-hydrogen atom positional and thermal parameters (at first isotropic, then anisotropic) followed. Hydrogen atoms were incorporated at their calculated positions in the later iterations which converged (max. shift/esd = 0.02) at $R = 0.050$ ($R_w = 0.063$). A final difference Fourier synthesis revealed no unusual features (max. 0.91, min. -0.84 e/Å³; both in the vicinity of the In atom). Crystallographic calculations were performed using the Enraf-Nonius Structure Determination Package.¹⁵ For all structure-factor calculations, neutral atom scattering factors and their anomalous dispersion corrections were taken from ref. 16. In the least-squares iterations, $\Sigma w\Delta^2$ [$w = 1/\sigma^2(|F_o|)$, $\Delta = (|F_o| - |F_c|)$] was minimized. Fractional atomic coordinates are listed in Table II; selected distances and angles are provided in Table III. A view of the solid-state structure of 1, with the atom numbering scheme, is presented in the Figure.

RESULTS AND DISCUSSION

From solution studies, the complexes $R_2\text{InOR}'$ where $R = \text{Me}$; $R' = \text{SiMe}_3$,⁴ SiPh_3 ,⁵ CMe_3 ,⁶ CPh_3 ,⁷ or $t\text{-Bu}$,¹⁰ $R = \text{Bu}$; $R' = \text{Bu}$ or Ph ¹¹ are postulated to exist primarily as dimers whereas when $R = \text{Me}$; $R' = \text{Me}$,⁸ $R = \text{CD}_3$; $R' = \text{Me}$ or CD_3 ,⁹ $R = \text{Et}$; $R' = \text{Me}$ or Et ⁹ the complexes are said to be trimeric. It is unfortunate that the limited solubility of 1 in organic solvents precludes the use of solution cryoscopic molecular weight determination. As is evident from the Figure, 1 exists in the solid state as a siloxy-bridged dimer. This degree of oligomerization is

undoubtedly due to size effects of the ligands bonded to the indium and oxygen atoms as the aforementioned trimeric alkoxides all involve relatively small substituents.

Though the formation of **1** was accomplished by a rather serendipitous event, this does not diminish the fundamental value that the elucidation of its structure and bonding geometry contributes to the general knowledge of alkoxide chemistry. As was stated earlier, the only previously reported structure of a simple organoindium alkoxide involved the dimer $(t\text{-Bu}_2\text{InOEt})_2$ **2**.¹² Interestingly, **2** was also prepared by a non-conventional method involving the reaction of $n\text{-BuLi}$, $\text{NHCMe}_2(\text{CH}_2)_3\text{CMe}_2$, and $t\text{-Bu}_2\text{InCl}$ in Et_2O solvent. The shortest intermolecular separations between the discrete dimeric units in crystals of **1** and **2** correspond to normal van der Waals distances. Compounds **1** and **2** both lie on crystallographic centres of symmetry and thus involve planar four-membered In_2O_2 cores. The mean In-O bond distances [2.154 Å **1**; 2.156 Å **2**] are essentially identical in both compounds, and the Si-O distance of 1.654(8) Å in **1** is in accord with the expected value of 1.63(2) Å.¹⁷ One of the oxygen atoms in **1** lies close to one of the phenyl ring planes [O-In-C(11)-C(12) torsion angle = 3(1)°] whereas both are nearly equidistant from the other phenyl ring plane [torsion angles: O-In-C(21)-C(26) = 46(1)°, O'-In-C(21)-C(26) = -39(1)°] which, accordingly, approximately bisects the O-In-O' angle. Ring aryl substituents are similarly oriented in crystals of $[\text{In}(\text{2,4,6-trimethylphenyl})_2\text{I}]_2$ **3**¹⁸ where the mean In-C distance of 2.169 Å associated with the bulkier ligands is slightly longer than that of 2.149 Å in **2**. In both **1** and **2**, the geometry about the oxygen atoms is distorted trigonal planar while that at the indium atoms is distorted tetrahedral. The O-In-O' and In-O-In' angles in **1** at 78.9 (3)° and 101.1 (3)°, respectively, are similar to corresponding values in **2** [75.2(3)° and 104.8 (2)°]. The considerable enlargement of the C-In-C angles [133.3(4)° in **1**, 127.2(4)° in **2**, 127.7(1)° in **3**] over the ideal

tetrahedral value is probably due principally to electronic factors with intramolecular non-bonded steric interactions between ring substituents also making some contribution.

SUPPLEMENTARY DATA

Additional material consisting of H-atom coordinates, thermal parameters, and a complete list of bond distances and angles are available from R.L.W.

ACKNOWLEDGEMENTS


We thank the Office of Naval Research for financial support.

REFERENCES

1. R. Nomura, S. Inazawa, H. Matsuda and S. Saeki, *Polyhedron*, **6**, 507 (1987).
2. R. Nomura, S. Fujii, K. Kanaya and H. Matsuda, *Polyhedron*, **9**, 361 (1990).
3. K. L. Chopra, S. Major and D. K. Pandya, *Thin Solid Films*, **102**, 1 (1983).
4. B. Armer and H. Schmidbaur, *Chem. Ber.*, **100**, 1521 (1967).
5. H. Schmidbaur, G. Kuhr and U. Kruger, *Angew. Chem. Int. Ed.*, **4**, 877 (1965).
6. H. Schmidbaur, *Angew. Chem. Int. Ed.*, **4**, 152 (1965).
7. H. Schmidbaur and F. Schindler, *Chem. Ber.*, **99**, 2178 (1966).
8. G. E. Coates and R. A. Whitcombe, *J. Chem. Soc.*, 3351(1956).
9. V. G. Man, H. Olapinski, R. Ott and J. Weidlein, *Z. Anorg. Allg. Chem.*, **410**, 195 (1974).
10. H. Schmidbaur, *Angew. Chem.* **77**, 169 (1965).
11. H. C. Clark and A. L. Pickard, *J. Organomet. Chem.*, **8**, 427 (1967).
12. D. C. Bradley, D. M. Frigo, M. B. Hursthouse and B. Hussain, *Organometallics*, **7**, 1112 (1988).
13. L. Chao and R. D. Rieke, *Syn. React. Inorg. Metal-Org. Chem.*, **4**, 373 (1974).
14. G. Becker and W. Holderich, *Chem. Ber.*, **108**, 2484 (1975).
15. Enraf-Nonius Structure Determination Package (SDP 3.0), Enraf-Nonius, Delft, The Netherlands (1985).
16. *International Tables for X-ray Crystallography*; The Kynoch Press, Birmingham, England (1974), Vol IV.
17. F. H. Allen, O. Kennard, D. G. Watson, L. Brammer, A. G. Orpen and R. Taylor, *J. Chem. Soc., Perkin Trans. 2*, S2 (1987).
18. J. T. Leman, J. W. Ziller and A. R. Barron, *Organometallics*, **10**, 1766 (1991).

Caption to Figure

Figure. ORTEP diagram (40% probability ellipsoids) showing the atom numbering scheme and solid-state conformation of $(\text{Ph}_2\text{InOSiMe}_3)_2$ **1**; primed atoms are related to the unprimed atoms by a crystallographic centre of symmetry. Small circles represent hydrogen atoms.



Figure

Table I
Crystallographic data for (Ph₂InOSiMe₃)₂ 1.

| | |
|--|--|
| Molecular formula | C ₃₀ H ₃₈ In ₂ O ₂ Si ₂ |
| Formula weight | 716.45 |
| Crystal system | monoclinic |
| Space group | <i>P</i> 2 ₁ / <i>c</i> (<i>C</i> _{2h} ⁵) - No.14 |
| <i>a</i> /Å | 9.136(1) |
| <i>b</i> /Å | 15.503(2) |
| <i>c</i> /Å | 11.826(1) |
| β(°) | 102.46(1) |
| No. of orientation refls.; θ(°) range | 25; 36-40 |
| <i>U</i> /Å ³ | 1635.5(6) |
| <i>Z</i> | 2 |
| <i>D_c</i> /g cm ⁻³ | 1.455 |
| μ/cm ⁻¹ (Cu- <i>K</i> α radiation, λ 1.5418 Å) | 124 |
| <i>T</i> /K | 298 |
| Crystal size/mm | 0.14 x 0.18 x 0.22 |
| <i>T</i> _{max} : <i>T</i> _{min} | 1.00:0.37 |
| Scan type | ω-2θ |
| Scanwidth/° | 1.00 + 0.14tanθ |
| θ _{max} /° | 75 |
| Intensity control refls. | 1 3 2, 1 1 3, 1 2 3, 2 3 1; |
| Overall variation in standards (%); | <2 |
| Repeat time /hr | 2 |
| Total no. of refls. (+ <i>h</i> , + <i>k</i> , ± <i>l</i>) recorded | 3590 |
| No. of non-equiv. refls. recorded | 3378 |
| <i>R</i> _{merge} (on <i>I</i>) | 0.038 |
| No. of refls. retained [<i>I</i> > 3.0σ(<i>I</i>)] | 1718 |
| No. of parameters refined | 163 |
| $R = \sum F_o - F_c / \sum F_o $ | 0.050 |
| $R_w = [\sum w(F_o - F_c)^2 / \sum w F_o ^2]^{1/2}$ | 0.063 |
| $S = [\sum w\Delta^2 / (N_{\text{obs.}} - N_{\text{para.}})]^{1/2}$ | 1.39 |
| Max. shift/esd in final least-squares cycle | 0.02 |
| Final Δρ(e/Å ³) max.;min. | 0.91 ; -0.84 |

Table II

Non-Hydrogen Atom Fractional Coordinates and equivalent isotropic thermal parameters ($\times 10^2$) for $(\text{Ph}_2\text{InOMe}_3)_2$ **1**, with Estimated Standard Deviations in Parentheses.

| Atom | <i>x</i> | <i>y</i> | <i>z</i> | <i>B</i> _{eq} (Å ²) |
|-------|-------------|------------|-------------|--|
| In | 0.03169(8) | 0.08616(5) | -0.07494(6) | 3.77(1) |
| Si | -0.2315(3) | 0.0991(2) | 0.0930(2) | 4.26(6) |
| O | -0.1169(7) | 0.0416(5) | 0.0320(5) | 3.7(1) |
| C(1) | -0.3412(13) | 0.1719(9) | -0.0152(10) | 6.6(3) |
| C(2) | -0.3531(13) | 0.0203(10) | 0.1476(10) | 6.3(4) |
| C(3) | -0.1190(16) | 0.1622(10) | 0.2162(11) | 7.1(4) |
| C(11) | -0.0888(12) | 0.0809(7) | -0.2525(6) | 4.2(2) |
| C(12) | -0.2364(12) | 0.0555(8) | -0.2769(9) | 4.7(3) |
| C(13) | -0.3160(15) | 0.0498(9) | -0.3938(11) | 6.3(3) |
| C(14) | -0.2451(16) | 0.0688(9) | -0.4801(11) | 7.2(4) |
| C(15) | -0.0961(19) | 0.0954(11) | -0.4547(9) | 9.1(4) |
| C(16) | -0.0218(13) | 0.1017(8) | -0.3413(10) | 5.4(3) |
| C(21) | 0.1857(10) | 0.1767(7) | 0.0214(9) | 3.9(2) |
| C(22) | 0.2155(14) | 0.2539(8) | -0.0280(9) | 5.5(3) |
| C(23) | 0.3158(15) | 0.3139(8) | 0.0392(13) | 6.9(4) |
| C(24) | 0.3860(13) | 0.2958(9) | 0.1477(11) | 6.3(3) |
| C(25) | 0.3569(14) | 0.2201(11) | 0.1967(11) | 6.8(4) |
| C(26) | 0.2568(14) | 0.1612(8) | 0.1327(10) | 5.7(3) |

Table III
Interatomic Distances (Å) and Angles (deg.) for (Ph₂InOMe₃)₂ 1,
with Estimated Standard Deviations in Parentheses.

| | | | |
|----------------|-----------|----------------|-----------|
| In-O | 2.159(7) | Si-O | 1.654(8) |
| In-O' | 2.149(8) | Si-C(1) | 1.834(12) |
| In-C(11) | 2.152(7) | Si-C(2) | 1.858(14) |
| In-C(21) | 2.135(10) | Si-C(3) | 1.868(13) |
| O-In-O' | 78.9(3) | C(1)-Si-C(3) | 110.4(6) |
| O-In-C(11) | 107.8(3) | C(2)-Si-C(3) | 110.1(6) |
| O-In-C(21) | 108.9(3) | In-O-In' | 101.1(3) |
| O'-In-C(11) | 106.2(3) | In-O-Si | 128.3(4) |
| O'-In-C(21) | 108.5(3) | In'-O-Si | 128.7(4) |
| C(11)-In-C(21) | 133.3(4) | In-C(11)-C(12) | 119.2(7) |
| O-Si-C(1) | 109.2(5) | In-C(11)-C(16) | 121.5(8) |
| O-Si-C(2) | 106.2(5) | In-C(21)-C(22) | 120.7(7) |
| O-Si-C(3) | 109.2(5) | In-C(21)-C(26) | 121.6(9) |
| C(1)-Si-C(2) | 111.7(6) | | |

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (2)*
Chemistry Division, Code 1113
800 North Quincy Street
Arlington, Virginia 22217-5000

Dr. James S. Murday (1)
Chemistry Division, Code 6100
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Robert Green, Director (1)
Chemistry Division, Code 385
Naval Air Weapons Center
Weapons Division
China Lake, CA 93555-6001

Dr. Elek Lindner (1)
Naval Command, Control and Ocean
Surveillance Center
RDT&E Division
San Diego, CA 92152-5000

Dr. Bernard E. Douda (1)
Crane Division
Naval Surface Warfare Center
Crane, Indiana 47522-5000

Dr. Richard W. Drisko (1)
Naval Civil Engineering
Laboratory
Code L52
Port Hueneme, CA 93043

Dr. Harold H. Singerman (1)
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Dr. Eugene C. Fischer (1)
Code 2840
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Defense Technical Information
Center (2)
Building 5, Cameron Station
Alexandria, VA 22314